

Utilisation of blood products at a centre in Pakistan. Is donating better than wasting?

Haya Ul Mujtaba¹, Nida Anwar², Naveena Fatima³, Imran Naseer⁴, Munira Borhany⁵, Tahir Shamsi⁶

Abstract

The study was designed to investigate the quantity and reasons of wastage of blood products. This was an observational study conducted from February 2018 to February 2019 at the National Institute of Blood Disease and Bone Marrow Transplantation (NIBD and BMT), PECHS campus. The study was approved by the institutional review board. Wastage and reasons of wastage for all the blood products were evaluated. Frequencies were calculated by using SPSS version 23.0. A total of 2,880 bags of blood products were available, including 960 each of platelets, packed red cells and fresh frozen plasma. The overall wastage rate was 3.5%. Packed red cells and platelets were fully consumed, yet shortage of supply was observed. However, highest wastage was observed in fresh frozen plasma i.e. 102 bags. Expiry of unused products 60 (59%) followed by broken bags 30 (29%) were two common modes of wastage. Wastage of blood products is a genuine issue in a hospital setup, strategies and plan of action should be discussed and implemented to ensure that they are available when and where they are needed most.

Keywords: Blood, blood component transfusion, blood bank.

DOI: <https://doi.org/10.47391/JPM.932>

Introduction

Blood is a precious resource of life¹ and one of the main life-saving interventions in various situations such as severe anaemia, trauma, surgery, chemotherapy, complicated pregnancy and childbirth.^{2,3} Most often blood products are needed urgently, and it is imminent to ensure the easy access.⁴ Moreover, there are many contributing factors for limited availability of blood such as lack of donors, underprivileged stock and transportation. According to the World Health Organization (WHO) about 87.5% of developing countries are unable to meet even 50% of the blood transfusion requirement of their population.⁵

A number of reasons are reported in literature for the wastage of blood products, including expired blood bags, broken seals, error in temperature maintenance, damaged bags, return of bags after dispatch, clotted and contaminated blood components.¹ In the United States, yearly wastage rate varies from 1% to 5% i.e. about 200,000 to one million units.⁶ However, Guyana, Iran and India reported the wastage rate of about 25%,⁷ 9.8%⁸ and 6.95% respectively.⁹

In Pakistan, about 1.2-1.5 million blood units are transfused annually.¹⁰ Recently it has been reported that Pakistan is facing a blood product shortage of about 40%.¹⁰ This shortage is followed by irrelevant use of 25% of whole blood and 80-85% of blood products.^{11,14} By focusing on the optimum use of blood and its products one can minimise the shortfall.

Limited studies have been carried out on the wastage rate of blood products in Pakistan. This study was conducted to evaluate the quantity and causes of wastage of blood products at the NIBD and BMT in order to minimise the wastage and review institutional policies to outsource blood products.

Materials and Methods

It was an observational study conducted at NIBD and BMT, PECHS campus Karachi, Pakistan from February 2018 - February 2019. Approval was taken from the institutional review board before starting the study. The study was based on statistical data available from the blood bank of the institute. Blood product consumption and wastage data was retrieved from internal software 'zaviya' which is used for maintaining a record of availability of blood product, issuance, usage and wastage. Variables for the study included total number of blood products made, number of each product, total number of wastage of blood products, number of wastage of each product and reasons of wastage. Microsoft Excel was used for preparation of the data sheet. Statistical analysis was done on Statistical Package for the Social Sciences (SPSS) version 23.0. Frequencies and percentages were calculated for qualitative variables.

Result


During the study period about 960 donations were

^{1,3,4}National Institute of Blood Disease and Bone Marrow Transplantation, Karachi, Pakistan; ^{2,5,6}Department of Haematology, National Institute of Blood Disease and Bone Marrow Transplantation, Karachi, Pakistan.

Correspondence: Nida Anwar. e-mail: drnidairfan@yahoo.com

Table: Reasons for wastage of blood products (n=102).

Reasons of wastage	n (%)
Expiry of unused products	60 (59)
Broken bags	30 (29)
Temperature Mismanagement	08 (8)
Contaminated products	04 (4)

**Figure :** Consumption and wastage rate for blood products.

received, from which 2,880 blood product bags were prepared including 960 (33.33%) each of PRBCs, PLT and FFP. Overall wastage rate for blood product was estimated as 102 (3.5%). PRBCs and PLT were fully consumed, although shortage was observed for both products. Out of 960 FFP only 858 (90%) were consumed and 102 (10%) were not consumed and were wasted. Consumption and wastage rates are shown in Figure 1.

The common reasons of wastage of blood products are listed in table 1.

Discussion

Blood is an important resource and its products are used in hospitals across the world to save human lives. Blood donations are limited, while its need is increasing globally and, hence, health care systems face lack of supply of blood products. Wastage of blood products is another important issue.⁸ This study was conducted to highlight the product consumption and wastage rate and also elaborate various reasons of wastage of blood products at NIBD and BMT in order to minimise the wastage and ensure the availability of blood products whenever needed. During the study, 960 donations were received from which 2,880 blood product bags were made, including 960 (33.33%) each of PRBCs, PLT and FFP. The overall wastage was estimated as 3.5%. PRBCs and PLT were fully consumed while the shortage of both products was seen which was fulfilled by arranging these products from outside. Out of the 960 (33.33%) FFP bags 102 (10%) were wasted due to various reasons, including the expiry of unused products 60 (59%), broken bags 30 (29%), temperature mismanagement 8 (8%) and contaminated products 4 (4%). A study conducted in

Guyana, from July 2012 to December 2014, reported the collection of 16,426 units of blood products. Blood wastage estimated for the years 2012, 2013 and 2014 were 487 (11.7%), 1,892 (45.5 %) and 1,788 (42.9 %) units respectively. According to Kurup R, et al reasons of wastage were broken bags 30 (1.7%), broken seal 99 (5.5%), broken cold chain 63 (3.5%), clotted blood 12 (0.7%), component with red cells 3 (0.2%), expired units 1,586 (87.7%) and return after 30 minutes 13 (0.7%).⁷ Similarly, our study revealed highest wastage due to expiry. In 2011, another study was conducted at Karnataka which reported the highest reason of wastage as blood contamination.¹⁵ In our study no such observation was found. In 2014, another study from Iran concluded the collection of 30,913 blood products including PRBCs 18,123 (58.6%), FFP 4,913 (15.8%), PLTs 6,695 (21.6%), whole blood 100 (0.3%) and cryoprecipitate 1,082 (3.5%) and reported the wastage rate of 9.85%. According to individual products, the wastage rate was 59.4% for PRBCs, 22% FFPs, 16% PLT and 2.4% cryoprecipitate.⁸ Similarly, a study conducted in India reported collection of 66,255 blood units out of which 44,617 were whole blood and other blood products were 21,638 PRBCs, 6,840 PLT, and 14,372 FFPs. The wastage rate was reported as 6.95%, out of which the wastage rate for PRBCs, PLTs, FFPs was 2.26%, 28.39% and 5.36% respectively.⁹ In our hospital, whole blood is not dispensed as per policy. In 2010, a study conducted at the teaching hospital of Semnan University in Iran revealed a high wastage of PRBCs which is contrary to our study findings. According to the study a total of 1,152 PRBCs were delivered to hospital out of which 759 (65.9%) were wasted and only 393 (34.1%) were consumed.¹⁶ According to WHO, some studies reported the high wastage reason for FFPs as short shelf life. Once prepared, FFPs have a shelf life of 24 hours.¹⁷ This could be the reason of FFPs expiry at our institute. Only 106 countries have national guidelines for blood transfusion. Hence, non-genuine transfusion is also an issue that leads to shortage of blood.⁸ In Pakistan, a study reported non-genuine transfusion rate in public and private sector as 45.90% and 30.60%.¹⁴ Australian Red Cross emphasised a number of policies that should be followed by blood banks and hospitals to reduce wastage rate including review temperature management plan, outsourcing of near expiry blood products, proper handling and good inventory system.¹⁸ World Health Organisation also recommends clinical guidelines for blood transfusion to be followed and prescribed when other techniques cannot viably control mortality. Availability of blood is limited. Hence, it is the necessity to execute policies for efficient transfusion and prevent wastage to fulfil blood requirement.¹⁷

Conclusion

In our study, highest wastage rate was observed for FFPs. However, PRBCs and PLT were fully consumed, shortage was faced for both products. There is a need to make policies and strategies to minimise wastage of blood products through collaboration with other blood banks. Donation is one of the options to utilise blood products more effectively, especially FFPs. There is also a need to educate people to donate blood voluntarily and also train the staff for better clinical use of blood products.

Disclaimer: Abstract has been presented at International society of blood transfusion (ISBT), Basel 2019 as poster.

Conflict of interest: None.

Funding disclosure: None

References

- Collins RA, Wisniewski MK, Waters JH, Triulzi DJ, Yazer MH. Effectiveness of multiple initiatives to reduce blood component wastage. *Am J Clin Pathol*. 2015; 143:329-35.
- Far RM, Rad FS, Abdolazimi Z, Kohan MMD. Determination of rate and causes of wastage of blood and blood products in Iranian hospitals. *Turk J Haematol*. 2014; 31:161-7.
- Javadzadeh Shahshahani H, Yavari MT, Attar M, Ahmadiyeh MH. Knowledge, Attitude and Practice (kap) Study about Blood Donation in the Urban Population of Yazd, Iran, 2004. *Trans Med*. 2006; 16:403-9.
- Bugge H, Karlsen N, Oydna E, Rake M, Wexels N, Bendabenda J, et al. A study of blood transfusion services at a district hospital in Malawi. *Vox sanguinis*. 2013; 104:37-45.
- Majdabadi HA, Kahouei M, Taslimi S, Langari M. Awareness of and attitude towards blood donation in students at the Semnan University of Medical Sciences. *Electron Physician*. 2018; 10:6821-8.
- Weatherall D, Clegg J. *Thalassemia in the Molecular Basis of Blood Disease*. 4th ed. Australia: Blackwell, 2001.
- Hannon T. Waste not, want not. *Am J Clin Pathol* 2015; 143:318-9.
- Kurup R, Anderson A, Boston C, Burns L, George M, Frank M. A study on blood product usage and wastage at the public hospital, Guyana. *BMC Res Notes*. 2016;9:307.
- Kanani AN, Vachhani JH, Dholakiya SK, Upadhyay SB. Analysis on discard of blood and its products with suggested possible strategies to reduce its occurrence in a blood bank of tertiary care hospital in Western India. *Global J Trans Med*. 2017; 2:130.
- World Health Organisation. Blood Safety: Situation analysis. [Online] [Cited 2008 January 25]. Available from: URL
- Horkey A. Thalassemia-A project for a General Pathology college class. [Online] [Cited 2012 February 10]. Available from: URL: <https://pathologyproject.wordpress.com/2012/02/10/thalassemia/>
- Ahmed P, Khan BA, Dilawar M, Mahmood A. Blood Transfusion: A Retrospective study at Combined Military Hospital, Multan. *Pak Armed Forces Med J*. 2000; 50: 76-80.
- Naeem MA, Masood A, Nuzhat M, Farhat AB. Evaluation of red cell transfusion triggers. *Pakistan J Pathol*. 2006; 17: 56-9.
- Afzal S. A comparison of public and private hospital on rational use of blood in Islamabad. *J Pak Med Assoc*. 2013; 63:85-9.
- Kora SA, Kulkarni K. An analysis of donor blood wastage in a blood bank in rural Karnataka. *J Clin Diagn Res*. 2011; 5:1393-6.
- Forouzesh F. Study of proper indications for blood transfusion in teaching hospital of Semnan University of Medical Sciences. 2010.
- WHO. Albanian strategy for safe blood transfusion. Tirana: WHO, 2010.
- Tool DINC, Calculator RC, Calculator RC. Health professionals. Victoria. 2012; 3: 0200.